U.S. DEPARTMENT OF COMMERCE
National Technical Information Service
PB80-144603

Distribution of Phytoplankton in Nevada Lakes

Environmental Monitoring and Support Lab, Las Vegas, NV

Dec 79

United States
Environmental Protection
Agency

Environmental Monitoring
and Support Laboratory
P.O. Box 15027
Las Vegas NV 89114

EPA-600/3-79-117
December 1979

PB80-144603

Research and Development

EPA

Distribution of Phytoplankton in Nevada Lakes

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

RESEARCH REPORTING SERIES

Research reports of the Office of Research and Development, U.S. Environmental Protection Agency, have been grouped into nine series. These nine broad categories were established to facilitate further development and application of environmental technology. Elimination of traditional grouping was consciously planned to foster technology transfer and maximum interface in related fields. The nine series are:

1. Environmental Health Effects Research
2. Environmental Protection Technology
3. Ecological Research
4. Environmental Monitoring
5. Socioeconomic Environmental Studies
6. Scientific and Technical Assessment Reports (STAR)
7. Interagency Energy—Environment Research and Development
8. "Special" Reports
9. Miscellaneous Reports

This report has been assigned to the ECOLOGICAL RESEARCH series. This series describes research on the effects of pollution on humans,plant and animal species, and materials. Problems are assessed for their long-and short-term influences. Investigations include formations, transport, and pathway studies to determine the fate of pollutants and their effects. This work provided the technical basis for setting standards to minimize undesirable changes in living organisms in the aquatic, terrestrial, and atmospheric environments.

This document is available to the public through the National Technical Information Service, Springfield, Virginia 22161

TECHNICAL REPORT DATA
(Please read Instructions on the reverse before completing)

1. REPORT NO.	2.	3. RECIPIENT'S ACCESSION NO.
EPA-600/3-79-117		PB 80 144603

4. TITLE AND SUBTITLE	5. REPORT DATE
DISTRIBUTION OF PHYTOPLANKTON IN NEVADA LAKES	December 1979
	6. PERFORMING ORGANIZATION CODE

7. AUTHOR(S)	8. PERFORMING ORGANIZATION REPORT NO.
V.W. Lambou, F.A. Morris, M.K. Morris, W.D. Taylor, L.R. Williams, and S.C. Hern	

9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT NO.
Environmental Monitoring and Support Laboratory Office of Research and Development U.S. Environmental Protection Agency Las Vegas, NV 89114	1BD884 11. CONTRACT/GRANT NO.

12. SPONSORING AGENCY NAME AND ADDRESS	13. TYPE OF REPORT AND PERIOD COVERED
U.S. Environmental Protection Agency-Las Vegas, NV Office of Research and Development Environmental Monitoring and Support Laboratory Las Vegas, NV 89114	02-21-75 to 12-11-75 14. SPONSORING AGENCY CODE EPA/600/07

15. SUPPLEMENTARY NOTES

16. ABSTRACT

This is a data report presenting the species and abundance of phytoplankton in the 10 lakes sampled by the National Eutrophication Survey in the State of Nevada. Results from the calculation of several water quality indices are also included (Nygaard's Trophic State Index, Palmer's Organic Pollution Index, and species diversity and abundance indices).

17. KEY WORDS AND DOCUMENT ANALYSIS

a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
*aquatic microbiology lakes *phytoplankton water quality	Nevada lake eutrophication Nygaard's trophic indices Palmer's organic pollution indices Species diversity and abundance	06 C, M 08 H 13 B

18. DISTRIBUTION STATEMENT	19. SECURITY CLASS (This Report)	21. NO. OF PAGES
RELEASE TO PUBLIC	UNCLASSIFIED	42
	20. SECURITY CLASS (This page) UNCLASSIFIED	22. PRICE

EPA Form 2220-1 (Rev. 4-77) PREVIOUS EDITION IS OBSOLETE

U.S. GOVERNMENT PRINTING OFFICE: 1979— 683-282/2220

EPA-600/3-79-117
December 1979

DISTRIBUTION OF PHYTOPLANKTON IN NEVADA LAKES

by

V. W. Lambou, F. A. Morris*, M. K. Morris*,
W. D. Taylor, L. R. Williams, and S. C. Hern

Water and Land Quality Branch
Monitoring Operations Division
Environmental Monitoring and Support Laboratory
Las Vegas, Nevada 89114

*Department of Biological Sciences
University of Nevada, Las Vegas
Las Vegas, Nevada 89154

ENVIRONMENTAL MONITORING AND SUPPORT LABORATORY
OFFICE OF RESEARCH AND DEVELOPMENT
U.S. ENVIRONMENTAL PROTECTION AGENCY
LAS VEGAS, NEVADA 89114

DISCLAIMER

This report has been reviewed by the Environmental Monitoring and Support Laboratory-Las Vegas, U.S. Environmental Protection Agency, and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

FOREWORD

Protection of the environment requires effective regulatory actions which are based on sound technical and scientific information. This information must include the quantitative description and linking of pollutant sources, transport mechanisms, interactions, and resulting effects on man and his environment. Because of the complexities involved, assessment of specific pollutants in the environment requires a total systems approach which transcends the media of air, water, and land. The Environmental Monitoring and Support Laboratory-Las Vegas contributes to the formation and enhancement of a sound monitoring data base for exposure assessment through programs designed to:

- develop and optimize systems and strategies for monitoring pollutants and their impact on the environment
- demonstrate new monitoring systems and technologies by applying them to fulfill special monitoring needs of the Agency's operating programs

This report presents the species and abundance of phytoplankton in the 10 lakes sampled by the National Eutrophication Survey in the State of Nevada, along with results from the calculation of several commonly used biological indices of water quality and community structure. These data can be used to biologically characterize the study lakes, and as baseline data for future investigations. This report was written for use by Federal, State, and local governmental agencies concerned with water quality analysis, monitoring, and/or regulation. Private industry and individuals similarly involved with the biological aspects of water quality will find the document useful. For further information contact the Water and Land Quality Branch, Monitoring Operations Division.

George B. Morgan
Director
Environmental Monitoring and Support Laboratory
Las Vegas

CONTENTS

	Page
Foreword	iii
Introduction	1
Materials and Methods	2
Lake and Site Selection	2
Sample Preparation	2
Examination	3
Quality Control	4
Results	5
Nygaard's Trophic State Indices	5
Palmer's Organic Pollution Indices	5
Species Diversity and Abundance Indices	7
Species Occurrence and Abundance	9
Literature Cited	10
Appendix A. Phytoplankton Species list for the State of Nevada	11
Appendix B. Summary of Phytoplankton Data	14

INTRODUCTION

The collection and analysis of phytoplankton data were included in the National Eutrophication Survey in an effort to determine relationships between algal characteristics and trophic status of individual lakes.

During spring, summer, and fall of 1975, the Survey sampled 156 lakes in 11 States. Over 450 algal species and varieties were identified and enumerated from the 430 water samples examined.

This report presents the species and abundance of phytoplankton in the 10 lakes sampled in the State of Nevada (Table 1). The Nygaard's Trophic State (Nygaard 1949), Palmer's Organic Pollution (Palmer 1969), and species diversity and abundance indices are also included.

TABLE 1. LAKES SAMPLED IN THE STATE OF NEVADA

STORET No.	Lake Name	County
3201	Lake Mead	Clark (Mohave in Ariz.)
3202	Lahontan Reservoir	Lyon, Churchill
3204	Rye Patch Reservoir	Pershing
3205	Lake Tahoe	Washoe, Carson City, Douglas (Placer, El Dorado in Calif.)
3206	Topaz Reservoir	Douglas (Mono in Calif.)
3207	Upper Pahranagat Lake	Lincoln
3208	Washoe Lake	Washoe
3209	Wildhorse Reservoir	Elko
3210	Wilson Sink Reservoir	Elko
3211	Walker Lake	Mineral

MATERIALS AND METHODS

LAKE AND SITE SELECTION

Lakes and reservoirs included in the Survey were selected through discussions with State water pollution agency personnel and U.S. Environmental Protection Agency Regional Offices (U.S. Environmental Protection Agency 1975). Screening and selection strongly emphasized lakes with actual or potential accelerated eutrophication problems. As a result, the selection was limited to lakes:

(1) impacted by one or more municipal sewage treatment plant outfalls either directly into the lake or by discharge to an inlet tributary within approximately 40 kilometers of the lake;

(2) 40 hectares or larger in size; and

(3) with a mean hydraulic retention time of at least 30 days.

Specific selection criteria were waived for some lakes of particular State interest.

Sampling sites for a lake were selected based on available information on lake morphometry, potential major sources of nutrient input, and on-site judgment of the field limnologist (U.S. Environmental Protection Agency 1975). Primary sampling sites were chosen to reflect the deepest portion of each major basin in a test lake. Where many basins were present, selection was guided by nutrient source information on hand. At each sampling site, a depth-integrated phytoplankton sample was taken. Depth-integrated samples were uniform mixtures of water from the surface to a depth of 15 feet (4.6 meters) or from the surface to the lower limit of the photic zone representing 1 percent of the incident light, whichever was greater. If the depth at the sampling site was less than 15 feet (4.6 meters), the sample was taken from just off the bottom to the surface. Normally, a lake was sampled three times in 1 year, providing information on spring, summer, and fall conditions.

SAMPLE PREPARATION

To preserve the sample 4 milliliters (ml) of Acid-Lugol's solution (Prescott 1970) were added to each 130-ml sample from each site at the time of collection. The samples were shipped to the Environmental Monitoring and Support Laboratory, Las Vegas, Nevada, where equal volumes from each site

were mixed to form two 130-ml composite samples for a given lake. One composite sample was put into storage and the other was used for the examination.

Prior to examination, the composite samples were concentrated by the settling method. Solids were allowed to settle for at least 24 hours prior to siphoning off the supernate. The volume of the removed supernate and the volume of the remaining concentrate were measured and concentrations determined. A small (8-ml) library subsample of the concentrate was then taken. The remaining concentrate was gently agitated to resuspend the plankton and poured into a capped, graduated test tube. If a preliminary examination of a sample indicated the need for a more concentrated sample, the contents of the test tube were further concentrated by repeating the settling method. Final concentrations varied from 15 to 40 times the original.

Permanent slides were prepared from concentrated samples after analysis was complete. A ring of clear Karo® corn syrup with phenol (a few crystals of phenol were added to each 100 ml of syrup) was placed on a glass slide. A drop of superconcentrate from the bottom of the test tube was placed in the ring. This solution was thoroughly mixed and topped with a coverglass. After the syrup at the edges of the coverglass had hardened, the excess was scraped away and the mount was sealed with clear fingernail polish. Permanent diatom slides were prepared by drying sample material on a coverglass, heating in a muffle furnace at 400° C for 45 minutes, and mounting in Hyrax®. Finally, the mounts were sealed with clear fingernail polish.

Backup samples, library samples, permanent sample slides, and Hyrax®-mounted diatom slides are being stored and maintained at the Environmental Monitoring and Support Laboratory-Las Vegas.

EXAMINATION

The phytoplankton samples were examined with the aid of binocular compound microscopes. A preliminary examination was performed to precisely identify and list all forms encountered. The length of this examination varied depending on the complexity of the sample. An attempt was made to find and identify all of the forms present in each sample. Often forms were observed which could not be identified to species or to genus. Abbreviated descriptions were used to keep a record of these forms (e.g., lunate cell, blue-green filament, Navicula #1). Diatom slides were examined using a standard light microscope. If greater resolution was essential to accurately identify the diatoms, a phase-contrast microscope was used.

After the species list was compiled, phytoplankton were enumerated using a Neubauer Counting Chamber with a 40X objective lens and a 10X ocular lens. All forms within each field were counted. The count was continued until a minimum of 100 fields had been viewed, or until the dominant form had been observed a minimum of 100 times.

®Registered trademark

QUALITY CONTROL

Project phycologists performed internal quality control intercomparisons regularly on 7 percent of the species identification and counts. Although an individual had primary responsibility for analyzing a sample, taxonomic problems were discussed among the phycologists.

Additional quality control checks were performed on the Survey samples by Dr. G. W. Prescott of the University of Montana at the rate of 5 percent. Quality control checks were made on 75 percent of these samples to verify species identifications while checks were made on the remaining 25 percent of the samples to verify genus counts. Presently, the agreement between quality control checks for species identification and genus enumerations is satisfactory.

RESULTS

A phytoplankton species list for the State is presented in Appendix A. Appendix B summarizes all of the phytoplankton data collected from the State by the Survey. The latter is organized by lake, and includes an alphabetical phytoplankton species list with concentrations for individual species given by sampling date. Results from the application of several indices are presented (Nygaard's Trophic State, Palmer's Organic Pollution, and species diversity and abundance). Each lake has been assigned a four-digit STORET number. (STORET (STOrage and RETrieval) is the U.S. Environmental Protection Agency's computer system which processes and maintains water quality data.) The first two digits of the STORET number identify the State; the last two digits identify the lake.

NYGAARD'S TROPHIC STATE INDICES

Five indices devised by Nygaard (1949) were proposed under the assumption that certain algal groups are indicative of levels of nutrient enrichment. These indices were calculated in order to aid in determining the surveyed lakes' trophic status. As a general rule, Cyanophyta, Euglenophyta, centric diatoms, and members of the Chlorococcales are found in waters that are eutrophic (rich in nutrients), while desmids and many pennate diatoms generally cannot tolerate high nutrient levels and so are found in oligotrophic waters (poor in nutrients).

In applying the indices to the Survey data, the number of taxa in each major group was determined from the species list for each sample. The ratios of these groups give numerical values which can be used as a biological index of water richness. The five indices and the ranges of values established for Danish lakes by Nygaard for each trophic state are presented in Table 2. The appropriate symbol, (E) eutrophic and (O) oligotrophic, follows each calculated value in the tables in Appendix B. A question mark (?) following a calculated value in these tables was entered when that value was within the range of both classifications.

PALMER'S ORGANIC POLLUTION INDICES

Palmer (1969) analyzed reports from 165 authors and developed algal pollution indices for use in rating water samples with high organic pollution. Two lists of organic-pollution-tolerant forms were prepared, one containing 20 genera, the other, 20 species (Tables 3 and 4). Each form was assigned a pollution index number ranging from 1 for moderately tolerant forms to 6 for

TABLE 2. NYGAARD'S TROPHIC STATE INDICES ADAPTED FROM HUTCHINSON (1967)

Index	Calculation	Oligotrophic	Eutrophic
Myxophycean	$\frac{\text{Myxophyceae}}{\text{Desmideae}}$	0.0-0.4	0.1-3.0
Chlorophycean	$\frac{\text{Chlorococcales}}{\text{Desmideae}}$	0.0-0.7	0.2-9.0
Diatom	$\frac{\text{Centric Diatoms}}{\text{Pennate Diatoms}}$	0.0-0.3	0.0-1.75
Euglenophyte	$\frac{\text{Euglenophyta}}{\text{Myxophyceae + Chlorococcales}}$	0.0-0.2	0.0-1.0
Compound	$\frac{\text{Myxophyceae + Chlorococcales + Centric Diatoms + Euglenophyta}}{\text{Desmideae}}$	0.0-1.0	1.2-25

TABLE 3. ALGAL GENUS POLLUTION INDEX (Palmer 1969)

Genus	Pollution Index
Anacystis	1
Ankistrodesmus	2
Chlamydomonas	4
Chlorella	3
Closterium	1
Cyclotella	1
Euglena	5
Gomphonema	1
Lepocinclis	1
Melosira	1
Micractinium	1
Navicula	3
Nitzschia	3
Oscillatoria	5
Pandorina	1
Phacus	2
Phormidium	1
Scenedesmus	4
Stigeoclonium	2
Synedra	2

TABLE 4. ALGAL SPECIES POLLUTION INDEX (Palmer 1969)

Species	Pollution Index
Ankistrodesmus falcatus	3
Arthrospira jenneri	2
Chlorella vulgaris	2
Cyclotella meneghiniana	2
Euglena gracilis	1
Euglena viridis	6
Gomphonema parvulum	1
Melosira varians	2
Navicula cryptocephala	1
Nitzschia acicularis	1
Nitzschia palea	5
Oscillatoria chlorina	2
Oscillatoria limosa	4
Oscillatoria princeps	1
Oscillatoria putrida	1
Oscillatoria tenuis	4
Pandorina morum	3
Scenedesmus quadricauda	4
Stigeoclonium tenue	3
Synedra ulna	3

extremely tolerant forms. Palmer based the index numbers on occurrence records and/or where emphasized by the authors as being especially tolerant of organic pollution.

In analyzing a water sample, any of the 20 genera or species of algae present in concentrations of 50 per milliliter or more are recorded. The pollution index numbers of the algae present are totaled, providing a genus score and a species score. Palmer determined that a score of 20 or more for either index can be taken as evidence of high organic pollution, while a score of 15 to 19 is taken as probable evidence of high organic pollution. Lower figures suggest that the organic pollution of the sample is not high, that the sample is not representative, or that some substance or factor interfering with algal persistence is present and active.

SPECIES DIVERSITY AND ABUNDANCE INDICES

"Information content" of biological samples is being used commonly by biologists as a measure of diversity. Diversity in this connection means the degree of uncertainty attached to the specific identity of any randomly selected individual. The greater the number of taxa and the more equal their proportions, the greater the uncertainty, and hence, the diversity (Pielou 1966). There are several methods of measuring diversity, e.g., the formulas given by Brillouin (1962) and Shannon and Weaver (1963). The method which is appropriate depends on the type of biological sample on hand.

Pielou (1966) classifies the types of biological samples and gives the measure of diversity appropriate for each type. The Survey phytoplankton samples are what she classifies as larger samples (collections in Pielou's terminology) from which random subsamples can be drawn. According to Pielou, the average diversity per individual (H) for these types of samples can be estimated from the Shannon-Wiener formula (Shannon and Weaver 1963):

$$H = -\sum_{i=1}^{S} P_i \log_x P_i$$

where P is the proportion of the ith taxon in the sample, which is calculated from n_i/N; n_i is the number of individuals per milliliter of the ith taxon; N is the total number of individuals per ml; and S is the total number of taxa. However, Basharin (1959) and Pielou (1966) have pointed out that H calculated from the subsample is a biased estimator of the sample H, and if this bias is to be accounted for, we must know the total number of taxa present in the sample since the magnitude of this bias depends on it.

Pielou (1966) suggests that if the number of taxa in the subsample falls only slightly short of the number in the larger sample, no appreciable error will result in considering S, estimated from the subsample, as being equal to the sample value. Even though considerable effort was made to find and identify all taxa, the Survey samples undoubtedly contain a fair number of rare phytoplankton taxa which were not encountered.

In the Shannon-Wiener formula, an increase in the number of taxa and/or an increase in the evenness of the distribution of individuals among taxa will increase the average diversity per individual from its minimal value of zero. Sager and Hasler (1969) found that the richness of taxa was of minor importance in determination of average diversity per individual for phytoplankton and they concluded that phytoplankton taxa in excess of the 10 to 15 most abundant ones have little effect on H. This was verified by our own calculations. Our counts are in number per milliliter and since logarithms to the base 2 were used in our calculations, H is expressed in units of bits per individual. When individuals of a taxon were so rare that they were not counted, a value of 1/130 per milliliter or 0.008 per milliliter was used in the calculations since at least one individual of the taxon must have been present in the collection.

A Survey sample for a given lake represents a composite of all phytoplankton collected at different sampling sites on the lake during a given sampling period. Since the number of samples (M) making up a composite is a function of both the complexity of the lake sampled and its size, it should affect the richness-of-taxa component of the diversity of our phytoplankton collections. The maximum diversity (MaxH) (i.e., when the individuals are distributed among the taxa as evenly as possible) was estimated from $\log_2 S$ (Pielou 1966), while the minimum diversity (MinH), was estimated from the formula:

$$\text{MinH} = -\frac{S-1}{N}\log_2\frac{1}{N} - \frac{N-(S-1)}{N}\log_2\frac{N-(S-1)}{N}$$

given by Zand (1976). The total diversity (D) was calculated from HN (Pielou 1966). Also given in Appendix B are L (the mean number of individuals per taxa per milliliter) and K (the number of individuals per milliliter of the most abundant taxon in the sample).

The evenness component of diversity (J) was estimated from H/MaxH (Pielou 1966). Relative evenness (RJ) was calculated from the formula:

$$\text{RJ} = \frac{\text{H-MinH}}{\text{MaxH-MinH}}$$

given by Zand (1976). Zand suggests that RJ be used as a substitute for both J and the redundancy expression given by Wilhm and Dorris (1968). As pointed out by Zand, the redundancy expression given by Wilhm and Dorris does not properly express what it is intended to show, i.e., the position of H in the range between MaxH and MinH. RJ may range from 0 to 1; being 1 for the most even samples and 0 for the least even samples.

Zand (1976) suggests that diversity indices be expressed in units of "sits", i.e., in logarithms to base S (where S is the total number of taxa in the sample) instead of in "bits", i.e., in logarithms to base 2. Zand points out that the diversity index in sits per individual is a normalized number ranging from 1 for the most evenly distributed samples to 0 for the least evenly distributed samples. Also, it can be used to compare different samples, independent of the number of taxa in each. The diversity in bits per

individual should not be used in direct comparisons involving various samples which have different numbers of taxa. Since MaxH equals log S, the expression in sits is equal to $\log_S S$, or 1. Therefore diversity in sits per individual is numerically equivalent to J, the evenness component for the Shannon-Wiener formula.

SPECIES OCCURRENCE AND ABUNDANCE

The alphabetic phytoplankton species list for each lake, presented in Appendix B, gives the concentrations of individual species by sampling date. Concentrations are in cells, colonies, or filaments (CEL, COL, FIL) per milliliter. An "X" after a species name indicates that the species identified in the preliminary examination was in such a low concentration that it did not appear in the count. A blank space indicates that the organism was not found in the sample collected on that date. Column S is used to designate the examiner's subjective opinion of the five dominant taxa in a sample, based upon relative size and concentration of the organism. The percent column (%C) presents, by abundance, the percentage composition of each taxon.

LITERATURE CITED

Basharin, G. P. 1959. On a statistical estimate for the entropy of a sequence of independent random variables, pp. 333-336. In: Theory of Probability and Its Applications (translation of "Teoriya Veroyatnosei i ee Premeneniya"). N. Artin (ed). 4. Society for Industrial and Applied Mathematics, Philadelphia.

Brillouin, L. 1962. Science and Information Theory (2nd ed.). Academic Press, New York. 351 pp.

Hutchinson, G. E. 1967. A Treatise on Limnology. II. Introduction to Lake Biology and the Limnoplankton. John Wiley and Sons, Inc., New York. 1,115 pp.

Nygaard, G. 1949. Hydrobiological studies of some Danish ponds and lakes. II. (K danske Vidensk. Selsk.) Biol. Sci. 7:293.

Palmer, C. M. 1969. A composite rating of algae tolerating organic pollution. J. Phycol. 5:78-82.

Pielou, E. C. 1966. The measurement of diversity in different types of biological collections. J. Theor. Biol. 13:131-144.

Prescott, G. W. 1970. How to Know the Freshwater Algae. William C. Brown Company, Dubuque. 348 pp.

Sager, P. E., and A. D. Hasler. 1969. Species diversity in lacustrine phytoplankton. I. The components of the index of diversity from Shannon's formula. Amer. Natur. 103(929):51-59.

Shannon, C. E., and W. Weaver. 1963. The Mathematical Theory of Communication. University of Illinois Press, Urbana. 117 pp.

U.S. Environmental Protection Agency. 1975. National Eutrophication Survey Methods 1973-1976. Working Paper No. 175. Environmental Monitoring and Support Laboratory, Las Vegas, Nevada, and Corvallis Environmental Research Laboratory, Corvallis, Oregon. 91 pp.

Wilhm, V. L., and T. C. Dorris. 1968. Biological parameters for water quality criteria. Bio-Science. 18:477.

Zand, S. M. 1976. Indexes associated with information theory in water quality. J. Water Pollut. Contr. Fed. 48(8):2026-2031.

APPENDIX A

PHYTOPLANKTON SPECIES LIST FOR THE STATE OF NEVADA

Actinastrum hantzschii
v. *fluviatile*
Anabaena
Ankistrodesmus falcatus
Ankistrodesmus falcatus
v. *mirabilis*
Aphanizomenon flos-aquae
Aphanothece
Asterionella formosa
Botryococcus braunii
Caloneis amphisbaena
Ceratium hirundinella
Ceratium hirundinella
f. *furcoides*
Chlamydomonas
Chroococcus
Closterium
Cocconeis pediculus
Cocconeis placentula
Coelastrum microporum
Coelastrum sphaericum
Cosmarium
Crucigenia quadrata
Crucigenia tetrapedia
Cryptomonas erosa
Crytomonas marssonii
Cyclotella meneghiniana
Cylindrotheca gracilis
Cylindrotheca spiralis
Cymatopleura solea
Cymbella affinis
Dactylococcopsis
Diatoma vulgare
Dictyosphaerium pulchellum
Dinobryon divergens
Entomoneis ornata
Epithemia turgida
Euglena acus
Eunotia pectinalis
v. *ventricosa*
Fragilaria brevistriata
Fragilaria brevistriata
v. *trigibba*
Fragilaria crotonensis
Glenodinium
Gloeocapsa
Gloeotrichia echinulata
Gomphonema olivaceum
Gomphonema truncatum
Gymnodinium ordinatum
Gyrosigma
Hantzschia amphioxys
Lyngbya
Melosira distans
Melosira granulata
Melosira granulata
v. *angustissima*
Melosira italica
Melosira varians
Merismopedia minima
Mesostigma viridis
Microcystis aeruginosa
Microcystis incerta
Mougeotia
Navicula exigua
Nitzschia filiformis
Nodularia
Oocystis
Opephora
Oscillatoria
Pandorina morum
Pediastrum boryanum
Pediastrum duplex
v. *reticulatum*
Phacus helikoides
Phormidium mucicola
Pinnularia
Quadrigula chodatii
Raphidiopsis curvata
Rhoicosphenia curvata
Rhopalodia gibba
Scenedesmus acuminatus
Scenedesmus dimorphus
Scenedesmus intermedius
v. *acaudatus*
Scenedesmus quadricauda
Schroederia setigera
Skeletonema potamos
Sphaerocystis schroeteri
Staurastrum
Stauroneis
Stephanodiscus
Surirella ovalis
Surirella ovata
Synedra ulna
Tabellaria fenestrata
Tetraedron minmum
Tetraedron minimum
v. *scrobiculatum*
Tetraedron muticum
Tetrastrum glabrum
Trachelomonas acanthostoma ?

Trachelomonas oblonga
v. *australica*
Trachelomonas volvocina

Trachelomonas volvocina
v. *punctata*

APPENDIX B. SUMMARY OF PHYTOPLANKTON DATA

This appendix was generated by computer. Because it was only possible to use upper case letters in the printout, all scientific names are printed in upper case and are not italicized.

The alphabetic phytoplankton lists include taxa without species names (e.g., EUNOTIA, EUNOTIA #1, FLAGELLATE, FLAGELLATES, MICROCYSTIS INCERTA ?, CHLOROPHYTAN COCCOID CELLED COLONY). When species determinations were not possible, symbols or descriptive phrases were used to separate taxa for enumeration purposes. Each name on a list, however, represents a unique species different from any other name on the same list, unless otherwise noted, for counting purposes.

Numbers were used to separate unidentified species of the same genus. A generic name listed alone is also a unique species. A question mark (?) is placed immediately after the portion of a name which was assigned with uncertainty. Numbered, questioned, or otherwise designated taxa were established on a lake-by-lake basis; therefore NAVICULA #2 from lake A cannot be compared to NAVICULA #2 from lake B. Pluralized categories (e.g., FLAGELLATES, CENTRIC DIATOMS, SPP.) were used for counting purposes when taxa could not be properly differentiated on the counting chamber.

LAKE NAME: LAKE MEAD
STORET NUMBER: 3201

NYGAARD TROPHIC STATE INDICES

DATE	02 24 75	06 11 75
MYXOPHYCEAN	01/0 E	0/01 O
CHLOROPHYCEAN	02/0 E	2,00 E
EUGLENOPHYTE	0/03 ?	0/02 ?
DIATOM	1,00 E	0,33 E
COMPOUND	04/0 E	3,00 E

PALMER'S ORGANIC POLLUTION INDICES

DATE	02 24 75	06 11 75
GENUS	00	00
SPECIES	00	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		02 24 75	06 11 75
AVERAGE DIVERSITY	H	1,35	0,87
NUMBER OF TAXA	A	8,00	11,00
NUMBER OF SAMPLES COMPOSITED	N	14,00	14,00
MAXIMUM DIVERSITY	MAXH	3,00	3,46
MINIMUM DIVERSITY	MINH	0,12	0,56
TOTAL DIVERSITY	D	812,70	133,98
TOTAL NUMBER OF INDIVIDUALS/ML	M	602,00	154,00
EVENNESS COMPONENT	J	0,45	0,25
RELATIVE EVENNESS	RJ	0,43	0,11
MEAN NUMBER OF INDIVIDUALS/TAXA	L	75,25	14,00
NUMBER/ML OF MOST ABUNDANT TAXON	K	417,00	110,00

LAKE NAME: LAKE MEAD
STORET NUMBER: 3201

CONTINUED

TAXA	FORM	02 24 78			06 11 78		
		1B	%C	ALGAL UNITS PER ML	1B	%C	ALGAL UNITS PER ML
CENTRIC DIATOM	CEL			X			
CERATIUM HIRUNDINELLA	CEL						X
CHROOMONAS ?	CEL	13	18,6	93			
CHROOMONAS ? ACUTA	CEL	11	69,3	417	11	71,4	110
CRYPTOMONAS	CEL						X
CRYPTOMONAS EROSA	CEL	19	7,6	46			
DIATOMA VULGARE	CEL						X
DINOBRION DIVERGENS	CEL						X
FRAGILARIA CROTONENSIS	CEL			X			X
MERISMOPEDIA MINIMA	COL			X			
OOCYSTIS	COL						X
RHOICOSPHENIA CURVATA	CEL				13	28,6	44
SCHROEDERIA SETIGERA	CEL			X			
SPHAEROCYSTIS SCHROETERI	COL						X
STAURASTRUM	CEL						X
STEPHANODISCUS	CEL						X
TETRAEDRON MUTICUM	CEL	14	7,6	46			
TOTAL				603			154

LAKE NAME: LAHONTAN RES.
STORET NUMBER: 3202

NYGAARD TROPHIC STATE INDICES

DATE	03 17 75	07 09 75	11 07 75
MYXOPHYCEAN	01/0 E	01/0 E	0/0 O
CHLOROPHYCEAN	0/0 O	01/0 E	01/0 E
EUGLENOPHYTE	2.00 E	0/02 ?	0/01 ?
DIATOM	0.33 E	1.00 E	0.40 E
COMPOUND	06/0 E	04/0 E	05/0 E

PALMER'S ORGANIC POLLUTION INDICES

DATE	03 17 75	07 09 75	11 07 75
GENUS	07	01	07
SPECIES	00	00	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		03 17 75	07 09 75	11 07 75
AVERAGE DIVERSITY	H	2.25	1.13	2.93
NUMBER OF TAXA	S	20.00	8.00	17.00
NUMBER OF SAMPLES COMPOSITED	N	4.00	4.00	4.00
MAXIMUM DIVERSITY	MAXH	4.32	3.00	4.09
MINIMUM DIVERSITY	MINH	0.12	0.02	0.16
TOTAL DIVERSITY	D	4461.75	5209.30	3395.87
TOTAL NUMBER OF INDIVIDUALS/ML	M	1983.00	4610.00	1159.00
EVENNESS COMPONENT	J	0.52	0.38	0.72
RELATIVE EVENNESS	RJ	0.51	0.38	0.71
MEAN NUMBER OF INDIVIDUALS/TAXA	L	99.15	576.25	68.18
NUMBER/ML OF MOST ABUNDANT TAXON	K	684.00	3646.00	316.00

LAKE NAME: LAMONTAN RES.
STORET NUMBER: 3303

CONTINUED

TAXA	FORM	03 17 75			07 09 75			11 07 75		
		IB	%C	ALGAL UNITS PER ML	IB	%C	ALGAL UNITS PER ML	IB	%C	ALGAL UNITS PER ML
ANABAENA	FIL						X			
ASTERIONELLA FORMOSA	CEL	1	1.7	34	14	1.8	69			X
CENTRIC DIATOM	CEL	11	34.8	686				13	6.1	189
CHLAMYDOMONAS	CEL	3	8.6	171						
CHLOROPHYTAN LOBATE CELL	CEL			X						
CHROOMONAS ? ACUTA	CEL	14	18.8	389	13	9.0	413	13	13.6	198
COCCONEIS PLACENTULA	CEL			X						X
CRYPTOMONAS EROSA	CEL	3	1.7	34				14	4.6	83
CYLINDROTHECA GRACILIS	CEL									X
CYMBELLA	CEL									X
DACTYLOCOCCOPSIS	CEL	1	3.6	68						
DIATOMA VULGARE	CEL			X						
EUGLENA	CEL	1	1.7	34						
FRAGILARIA CROTONENSIS	CEL				11	79.1	3566			X
GLENODINIUM	CEL						X			
MELOSIRA #1	CEL			X						
MELOSIRA DISTANS	CEL							3	9.1	166
MELOSIRA GRANULATA	CEL				13	7.9	344			
MELOSIRA GRANULATA V. ANGUSTISSIMA	CEL				1	1.8	69			
MELOSIRA VARIANS	CEL			X						
NAVICULA	CEL							1	13.6	189
NITZSCHIA #1	CEL			X						
NITZSCHIA #2	CEL			X						
NITZSCHIA #3	CEL							11	27.3	316
NITZSCHIA SPP.	CEL	13	33.8	650						
ROICOSPHENIA CURVATA	CEL			X						X
SCHROEDERIA SETIGERA	CEL				1	1.8	69			
SKELETONEMA POTAMOS	CEL							1	9.1	108
STEPHANODISCUS	CEL									X
SURIRELLA OVATA	CEL			X				1	4.6	83
SYNEDRA ULNA	CEL			X						
TABELLARIA FENESTRATA	CEL			X				3	4.6	83
TETRAEDRON MINIMUM	CEL									X
TRACHELOMONAS	CEL			X						
TOTAL				1983			4619			1189

LAKE NAME: RYE PATCH RES.
STORET NUMBER: 3204

NYGAARD TROPHIC STATE INDICES

DATE	06 01 75	07 11 75	11 07 75
MYXOPHYCEAN	01/0 E	1.00 E	01/0 E
CHLOROPHYCEAN	0/0 O	1.00 E	04/0 E
EUGLENOPHYTE	1.00 E	0.50 E	0.20 ?
DIATOM	0.50 E	0.50 E	0.67 E
COMPOUND	04/0 E	4.00 E	08/0 E

PALMER'S ORGANIC POLLUTION INDICES

DATE	06 01 75	07 11 75	11 07 75
GENUS	00	00	05
SPECIES	00	00	03

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		06 01 75	07 11 75	11 07 75
AVERAGE DIVERSITY	H	0.02	1.01	2.35
NUMBER OF TAXA	S	8.00	8.00	15.00
NUMBER OF SAMPLES COMPOSITED	N	2.00	5.00	9.00
MAXIMUM DIVERSITY	MAXH	3.00	3.00	3.91
MINIMUM DIVERSITY	MINH	1.41	0.76	0.07
TOTAL DIVERSITY	D	0.60	68.68	6457.80
TOTAL NUMBER OF INDIVIDUALS/ML	N	30.00	68.00	2748.00
EVENNESS COMPONENT	J	0.01	0.34	0.60
RELATIVE EVENNESS	RJ	-0.87	0.12	0.60
MEAN NUMBER OF INDIVIDUALS/TAXA	L	3.75	8.50	183.20
NUMBER/ML OF MOST ABUNDANT TAXON	K	30.00	34.00	889.00

LAKE NAME: RYE PATCH RES.
STORET NUMBER: 3304

CONTINUED

TAXA	FORM	06 01 78			07 11 78			11 07 78		
		%	QC	ALGAL UNITS PER ML	%	QC	ALGAL UNITS PER ML	%	QC	ALGAL UNITS PER ML
ANABAENA	FIL			X			X			
CENTRIC DIATOM	CEL	11	100.	30						
CHLAMYDOMONAS	CEL							14	20.0	866
CHROOMONAS ? ACUTA	CEL							11	32.4	809
CLOSTERIUM	CEL						X			
CRYPTOMONAS EROSA	CEL				11	80.0	34		2.9	81
CYANOPHYTAN FILAMENT	FIL									X
CYCLOTELLA	CEL									X
CYMATOPLEURA	CEL			X						
CYMBELLA	CEL			X						
EUGLENA	CEL			X						
GOMPHONEMA	CEL			1						
GYROSIGMA	CEL						X			
MELOSIRA GRANULATA	CEL			X			X			X
NAVICULA	CEL									X
NITZSCHIA	CEL				23	80.0	34			
OOCYSTIS	COL							18	0.8	343
PANDORINA MORUM	COL							23	5.9	163
PENNATE DIATOM	CEL						X			X
PHACUS	CEL									
RHOPALODIA GIBBA	CEL			X						
SCENEDESMUS QUADRICAUDA	COL									X
SCHROEDERIA SETIGERA	CEL							23	36.8	727
SPHAEROCYSTIS SCHROETERI	COL						X			
SURIRELLA OVATA	CEL								2.9	81
TETRASTRUM GLABRUM	COL									X
TRACHELOMONAS ACANTHOSTOMA ?	CEL									X
TOTAL				30			68			3748

LAKE NAME: LAKE TAHOE
STORET NUMBER: 3206

NYGAARD TROPHIC STATE INDICES

DATE	03 18 75	07 02 75	11 04 75
MYXOPHYCEAN	0/01 O	0/01 O	0/01 O
CHLOROPHYCEAN	0/01 O	0/01 O	0/01 O
EUGLENOPHYTE	0/0 ?	01/0 E	0/0 ?
DIATOM	2,00 E	1,00 E	0,50 E
COMPOUND	2,00 E	2,00 E	1,00 O

PALMER'S ORGANIC POLLUTION INDICES

DATE	03 18 75	07 02 75	11 04 75
GENUS	00	03	00
SPECIES	00	00	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		03 18 75	07 02 75	11 04 75
AVERAGE DIVERSITY	H	1,00	0,00	1,58
NUMBER OF TAXA	S	5,00	4,00	5,00
NUMBER OF SAMPLES COMPOSITED	M	10,00	8,00	10,00
MAXIMUM DIVERSITY	MAXH	2,32	2,00	2,32
MINIMUM DIVERSITY	MINH	0,43	0,10	1,75
TOTAL DIVERSITY	D	70,00	0,00	12,64
TOTAL NUMBER OF INDIVIDUALS/ML	N	70,00	302,00	8,00
EVENNESS COMPONENT	J	0,43	0,00	0,68
RELATIVE EVENNESS	RJ	0,31	-0,05	-0,29
MEAN NUMBER OF INDIVIDUALS/TAXA	L	14,00	75,50	1,60
NUMBER/ML OF MOST ABUNDANT TAXON	K	35,00	302,00	3,00

LAKE NAME: LAKE TAHOE CONTINUED
STORET NUMBER: 3205

TAXA	FORM	03 10 75			07 02 75			11 04 75		
		1S	%C	ALGAL UNITS PER ML	1S	%C	ALGAL UNITS PER ML	1S	%C	ALGAL UNITS PER ML
CENTRIC DIATOM	CEL							137,01		3
CHROOMONAS ? ACUTA	CEL			X						
COSMARIUM	CEL			X			X			X
CYMBELLA	CEL							137,01		3
GYMNODINIUM ORDINATUM	CEL							128,01		3
MELOSIRA ITALICA	CEL			X						
NITZSCHIA	CEL	1150,01		35	1100,1		302			
SYNEDRA ULNA	CEL									X
TRACHELOMONAS	CEL						X			
TOTAL				70			302			8

LAKE NAME: TUPAI RES.
STORET NUMBER: 3206

NYGAARD TROPHIC STATE INDICES

DATE	03 19 75	06 30 75	11 05 75
MYXOPHYCEAN	0/01 O	04/0 E	2,00 E
CHLOROPHYCEAN	2,00 E	02/0 E	2,00 E
EUGLENOPHYTE	0/02 ?	0/06 ?	0/04 ?
DIATOM	1,00 E	0,40 E	2,00 E
COMPOUND	5,00 E	08/0 E	6,00 E

PALMER'S ORGANIC POLLUTION INDICES

DATE	03 19 75	06 30 75	11 05 75
GENUS	01	06	01
SPECIES	00	00	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		03 19 75	06 30 75	11 05 75
AVERAGE DIVERSITY	H	2,44	1,29	2,29
NUMBER OF TAXA	S	11,00	17,00	11,00
NUMBER OF SAMPLES COMPOSITED	N	2,00	2,00	2,00
MAXIMUM DIVERSITY	MAXH	3,46	4,09	3,46
MINIMUM DIVERSITY	MINH	0,15	0,02	0,19
TOTAL DIVERSITY	D	1742,16	16829,34	1284,69
TOTAL NUMBER OF INDIVIDUALS/ML	N	714,00	13046,00	561,00
EVENNESS COMPONENT	J	0,71	0,32	0,66
RELATIVE EVENNESS	RJ	0,70	0,32	0,65
MEAN NUMBER OF INDIVIDUALS/TAXA	L	64,91	767,41	51,00
NUMBER/ML OF MOST ABUNDANT TAXON	K	198,00	9825,00	224,00

LAKE NAME: TOPAZ RES.
STORET NUMBER: 3306

CONTINUED

TAXA	FORM	03 19 75			06 30 75			11 08 75		
		IB	%C	ALGAL UNITS PER ML	IB	%C	ALGAL UNITS PER ML	IB	%C	ALGAL UNITS PER ML
ANABAENA	FIL						X			X
APHANIZOMENON FLOS-AQUAE	FIL				13	0.2	01075	19	3.4	19
ASTERIONELLA FORMOSA	CEL			X	13	9.1	01191			
CENTRIC DIATOM #1	CEL	18	37.7	198						
CERATIUM HIRUNDINELLA	CEL									X
CHROOMONAS ? ACUTA	CEL		11.1	79	15	5.0	00655		3.4	19
COCCONEIS	CEL			X						
CRYPTOMONAS	CEL	4	5.6	40	15	0.2	30	13	39.9	224
CRYPTOMONAS EROSA	CEL						X			
CRYPTOMONAS MARSSONII	CEL						X			
DIATOMA VULGARE	CEL						X			
DICTYOSPHAERIUM PULCHELLUM	COL	11	16.7	119						
EPITHEMIA	CEL						X			
FRAGILARIA CROTONENSIS	CEL	12	22.3	159	11	75.3	09825	13	20.0	112
MELOSIRA	CEL	13	16.7	119	14	1.2	160	8	10.0	56
OOCYSTIS	COL						X			
OSCILLATORIA	FIL						00110			
PHORMIDIUM MUCICOLA	FIL						X			
SCHROEDERIA SETIGERA	CEL			X						X
SPHAEROCYSTIS SCHROETERI	COL						X			X
STAURASTRUM	CEL			X				11	20.0	112
STEPHANODISCUS	CEL			X			X	4	3.4	19
SURIRELLA OVATA	CEL						X			
TOTAL				714			13046			861

LAKE NAME: UPPER PAHRANAGAT LAKE
STORET NUMBER: 3207

NYGAARD TROPHIC STATE INDICES

DATE	05 06 75	08 21 75	11 21 75
MYXOPHYCEAN	0/0 D	03/0 E	2.00 E
CHLOROPHYCEAN	04/0 E	13/0 E	3.00 E
EUGLENOPHYTE	0.25 E	0.25 E	0.40 E
DIATOM	0/03 ?	0.14 ?	0.12 ?
COMPOUND	05/0 E	22/0 E	9.00 E

PALMER'S ORGANIC POLLUTION INDICES

DATE	05 06 75	08 21 75	11 21 75
GENUS	02	15	12
SPECIES	03	05	02

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		05 06 75	08 21 75	11 21 75
AVERAGE DIVERSITY	H	2.13	0.74	2.85
NUMBER OF TAXA	S	11.00	39.00	28.00
NUMBER OF SAMPLES COMPOSITED	M	3.00	2.00	2.00
MAXIMUM DIVERSITY	MAXH	3.46	5.29	4.81
MINIMUM DIVERSITY	MINH	0.12	0.02	0.13
TOTAL DIVERSITY	D	2017.11	30936.44	7623.75
TOTAL NUMBER OF INDIVIDUALS/ML	N	947.00	41806.00	2675.00
EVENNESS COMPONENT	J	0.62	0.14	0.59
RELATIVE EVENNESS	RJ	0.61	0.14	0.59
MEAN NUMBER OF INDIVIDUALS/TAXA	L	86.09	1071.95	95.54
NUMBER/ML OF MOST ABUNDANT TAXON	K	425.00	37834.00	899.00

LAKE NAME: UPPER PANRAGAT LAKE CONTINUED
STORET NUMBER: 3307

TAXA	FORM	09 06 75			09 21 75			11 21 75		
		IS	%C	ALGAL UNITS PER ML	IS	%C	ALGAL UNITS PER ML	IS	%C	ALGAL UNITS PER ML
ACTINASTRUM HANTZSCHII V. FLUVIATILE	COL					0.3	63			
ANKISTRODESMUS FALCATUS	CEL	15	6.9	65						
ANKISTRODESMUS FALCATUS V. MIRABILIS	CEL					0.9	378			
BOTRYOCOCCUS BRAUNII	COL						X			
CALONEIS AMPHISBAENA	CEL						X		0.9	33
CERATIUM HIRUNDINELLA	CEL						X			
CERATIUM HIRUNDINELLA F. FURCOIDES	CEL						X			
CHROOMONAS ? ACUTA	CEL	12	44.9	435					9.8	384
COCCONEIS	CEL						X			
COSMARIUM	CEL									X
CRUCIGENIA QUADRATA	COL					0.6	252			
CRUCIGENIA TETRAPEDIA	COL						X			
CRYPTOMONAS	CEL					0.3	63			
CRYPTOMONAS EROSA	CEL			X						
CRYPTOMONAS MARSSONII	CEL	11	17.2	163						
CYCLOTELLA MENEGHINIANA	CEL				14	2.1	903		8.2	139
CYLINDROTHECA GRACILIS	CEL							13	8.6	331
CYNATOPLEURA #1	CEL						X			
CYNATOPLEURA SOLEA	CEL						X			X
CYMBELLA	CEL									X
CYMBELLA SPP.	CEL						X			
DACTYLOCOCCOPSIS	CEL								1.7	66
EPITHEMIA	CEL						X			
EUGLENA	CEL				13	2.3	946			X
GYMNOZIGMA	CEL						X			X
MELOSIRA VARIANS	CEL						X		0.9	33
MERISMOPEDIA MINIMA	COL				13	90.8	37834			
MESOTIGMA VIRIDIS	CEL						X		0.9	33
MICROCYSTIS AERUGINOSA	COL						X			
MICROCYSTIS INCERTA	COL					0.3	63			
NAVICULA #1	CEL									X
NAVICULA #2	CEL							13	23.3	623
NITZSCHIA	CEL									X
NITZSCHIA SPP.	CEL							14	8.6	231
OOCYSTIS	CEL					0.6	252			
OSCILLATORIA	FIL								2.6	69
PEDIASTRUM BORYANUM	COL						X			X
PEDIASTRUM DUPLEX V. RETICULATUM	COL								0.9	33
PENNATE DIATOMS	CEL									
PHACUS HELIKOIDES	CEL				15	1.6	588	11	33.6	899
PINNULARIA #1	CEL				12	0.6	252			
PINNULARIA #2	CEL									X
QUADRIGULA CHODATII	CEL						X			
RHODOSPHENIA CURVATA	CEL	14	6.9	65						
RHOPALODIA GIBBA	CEL						X			
SCENEDESMUS ACUMINATUS	COL			X			X			
SCENEDESMUS DIMORPHUS	COL									X
SCENEDESMUS INTERMEDIUS V. ACAUDATUS	COL						X			
SCENEDESMUS QUADRICAUDA	COL					0.6	252			
SCHROEDERIA SETIGERA	CEL	13	20.7	196			X			X
SPHAEROCYSTIS SCHROETERI	COL		3.8	33			X			
SURIRELLA	CEL			X						
SURIRELLA OVALIS	CEL									
SURIRELLA OVATA	CEL									X
SYNEDRA	CEL						X	15	1.7	66
SYNEDRA ULNA	CEL			X			X		1.7	66
TETRAEDRON MINIMUM V. SCROBICULATUM	CEL						X			
TETRASTRUM GLABRUM	COL						X			
TRACHELOMONAS	CEL						X			
TRACHELOMONAS OBLONGA V. AUSTRALICA	CEL			X						
TRACHELOMONAS VOLVOCINA	CEL						X			X
TOTAL				947			41806			3675

LAKE NAME: WASHOE LAKE
STORET NUMBER: 3209

NYGAARD TROPHIC STATE INDICES

DATE	03 17 75	06 27 75	11 11 75
MYXOPHYCEAN	1.00 E	02/0 E	0/0 O
CHLOROPHYCEAN	2.00 E	01/0 E	01/0 E
EUGLENOPHYTE	0.67 E	0.67 E	0/01 ?
DIATOM	0.23 ?	0.18 ?	0.80 E
COMPOUND	8.00 E	07/0 E	06/0 E

PALMER'S ORGANIC POLLUTION INDICES

DATE	03 17 75	06 27 75	11 11 75
GENUS	01	07	01
SPECIES	00	00	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		03 17 75	06 27 75	11 11 75
AVERAGE DIVERSITY	H	1.89	2.59	2.39
NUMBER OF TAXA	S	25.00	19.00	16.00
NUMBER OF SAMPLES COMPOSITED	N	3.00	3.00	3.00
MAXIMUM DIVERSITY	MAXH	4.64	4.25	4.00
MINIMUM DIVERSITY	MINH	0.30	0.13	0.11
TOTAL DIVERSITY	D	1716.12	4330.48	4029.54
TOTAL NUMBER OF INDIVIDUALS/ML	M	908.00	1672.00	1686.00
EVENNESS COMPONENT	J	0.41	0.61	0.60
RELATIVE EVENNESS	RJ	0.37	0.60	0.59
MEAN NUMBER OF INDIVIDUALS/TAXA	L	36.32	88.00	105.38
NUMBER/ML OF MOST ABUNDANT TAXON	K	474.00	859.00	881.00

LAKE NAME: WASHOE LAKE
STORET NUMBER: 3208

CONTINUED

TAXA	FORM	03 17 75			06 27 75			11 11 75		
		1S	%C	ALGAL UNITS PER ML	1S	%C	ALGAL UNITS PER ML	1S	%C	ALGAL UNITS PER ML
CHROOMONAS ? ACUTA	CEL	151	8.7	79						
CLOSTERIUM	CEL			X						
COCCONEIS	CEL								4.6	77
COCCONEIS PLACENTULA	CEL				2.7	45				X
COELASTRUM SPHAERICUM	COL									
CRYPTOMONAS EROSA	CEL	141	8.7	79	151	9.1	136			X
CYCLOTELLA	CEL									
CYMATOPLEURA	CEL			X						X
CYMBELLA	CEL			X			X			
DACTYLOCOCCOPSIS	CEL			X						
ENTOMONEIS ORNATA	CEL			X						
EPITHEMIA	CEL							121	6.9	115
EPITHEMIA 01	CEL			X						
EPITHEMIA 02	CEL				141	2.7	45			
EUGLENA ACUS	CEL				121	8.1	136			X
EUNOTIA	CEL									
FRAGILARIA	CEL					5.4	90		6.9	115
GLOEOCAPSA	COL					2.7	45			
GOMPHONEMA	CEL									X
GOMPHONEMA 01	CEL			X						
GOMPHONEMA 02	CEL						X			
GOMPHONEMA TRUNCATUM	CEL			X						
HANTZSCHIA	CEL								2.3	38
HANTZSCHIA AMPHIOXYS	CEL			X						
MELOSIRA GRANULATA	CEL	121	21.7	197			X	111	52.3	891
MELOSIRA GRANULATA V. ANGUSTISSIMA	CEL	111	52.2	474	111	51.4	859	1	13.6	930
MELOSIRA ITALICA	CEL							131	4.6	77
MELOSIRA VARIANS	CEL			X						
MICROCYSTIS INCERTA	COL				2.7	45				
NAVICULA 01	CEL			X						
NAVICULA 02	CEL						X			
NITZSCHIA PTLIFORMIS	CEL									X
OOCYSTIS	COL			X						
PHACUS	CEL			X						
PINNULARIA	CEL			X	2.7	45				
RHOICOSPHENIA CURVATA	CEL					X				
RHOPALODIA GIBBA	CEL	131	8.7	79	131	9.1	136			
SCHROEDERIA SETIGERA	CEL			X		X				
STAURONEIS	CEL							151	4.6	77
STEPHANODISCUS	CEL							1	2.3	38
SURIRELLA	CEL							141	2.3	38
SURIRELLA 01	CEL			X						
SURIRELLA OVATA	CEL			X	2.7	45				
SYNEDRA ULNA	CEL			X	2.7	45				
TABELLARIA FENESTRATA	CEL			X						
TRACHELOMONAS OBLONGA V. AUSTRALICA	CEL			X						
TRACHELOMONAS VOLVOCINA	CEL						X			
TOTAL				900			1672			1696

LAKE NAME: WILDHORSE RES,
STORET NUMBER: 3209

WYGAARD TROPHIC STATE INDICES

DATE	05 30 75	08 01 75	11 06 75
MYXOPHYCEAN	03/0 E	04/0 E	02/0 E
CHLOROPHYCEAN	01/0 E	0/0 O	01/0 E
EUGLENOPHYTE	0,50 E	0/04 ?	0,33 E
DIATOM	0,14 ?	0,11 ?	0/02 ?
COMPOUND	07/0 E	05/0 E	04/0 E

PALMER'S ORGANIC POLLUTION INDICES

DATE	05 30 75	08 01 75	11 06 75
GENUS	04	01	00
SPECIES	00	00	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		05 30 75	08 01 75	11 06 75
AVERAGE DIVERSITY	H	0,60	2,55	1,45
NUMBER OF TAXA	S	16,00	15,00	7,00
NUMBER OF SAMPLES COMPOSITED	N	5,00	1,00	5,00
MAXIMUM DIVERSITY	MAXH	4,00	3,91	2,81
MINIMUM DIVERSITY	MINH	0,02	0,10	0,20
TOTAL DIVERSITY	D	5983,00	4365,60	427,75
TOTAL NUMBER OF INDIVIDUALS/ML	M	9805,00	1712,00	295,00
EVENNESS COMPONENT	J	0,15	0,65	0,52
RELATIVE EVENNESS	RJ	0,15	0,65	0,48
MEAN NUMBER OF INDIVIDUALS/TAXA	L	612,81	114,13	42,14
NUMBER/ML OF MOST ABUNDANT TAXON	K	8959,00	541,00	196,00

LAKE NAME: WILDHORSE RES, CONTINUED
STORET NUMBER: 3209

TAXA	FORM	05 30 78 IB	05 30 78 %C	05 30 78 ALGAL UNITS PER ML	08 01 78 IB	08 01 78 %C	08 01 78 ALGAL UNITS PER ML	11 06 78 IB	11 06 78 %C	11 06 78 ALGAL UNITS PER ML
APHANIZOMENON FLOS-AQUAE	FIL				13	15,0	270	11	11,3	33
APHANOTHECE	COL				13	5,3	90			
CENTRIC DIATOM	CEL	11	91,4	8950						
CHROOMONAS ? ACUTA	CEL							12	66,4	196
CYCLOTELLA BODANICA	CEL									X
CYCLOTELLA COMTA	CEL									X
CYCLOTELLA KUTZINGIANA	CEL									X
CYMBELLA TUMIDA	CEL									X
DINOBRYON SERTULARIA	CEL									X
EUNOTIA PECTINALIS	CEL									X
FRAGILARIA CROTONENSIS	CEL									X
GLOEOCYSTIS GIGAS	COL									X
GLOEOCYSTIS VASTA	COL									X
GOMPHONEMA ACUMINATUM	CEL									X
GOMPHONEMA OLIVACEUM	CEL									X
GYROSIGMA ACUMINATUM	CEL									X
MELOSIRA GRANULATA	CEL									X
MELOSIRA GRANULATA VAR. ANGUSTISSIMA	CEL									X
MELOSIRA ITALICA SUBSP. SUBARCTICA	CEL									X
MELOSIRA VARIANS	CEL									X
NAVICULA CRYPTOCEPHALA	CEL									X
NAVICULA EXIGUA	CEL									X
NAVICULA RADIOSA	CEL									X
NITZSCHIA ACICULARIS	CEL									X
NITZSCHIA AMPHIBIA	CEL									X
NITZSCHIA DISSIPATA	CEL									X
NITZSCHIA FILIFORMIS	CEL		0,4	40		2,6	45			
NITZSCHIA PALEA	CEL									X
OSCILLATORIA LIMOSA	FIL									X
PHORMIDIUM AUTUMNALE	FIL									X
PINNULARIA BRAUNII	CEL									X
PINNULARIA BRAUNII VAR. AMERICANA	CEL									X
PINNULARIA BRAUNII VAR. COMMUTATA	CEL									X
PINNULARIA BRAUNII VAR. GRUNOWII	CEL									X
PINNULARIA BRAUNII VAR. INTERRUPTA	CEL									X
PINNULARIA BRAUNII VAR. LINEARIS	CEL									X
PINNULARIA BRAUNII VAR. SUBCAPITATA	CEL									X
PINNULARIA BRAUNII VAR. UNDULATA	CEL									X
PINNULARIA BRAUNII VAR. VIRIDIS	CEL									X
PINNULARIA BREBISSONII	CEL									X
PINNULARIA BREBISSONII VAR. KRAMMERI	CEL									X
PINNULARIA BREBISSONII VAR. LANCEOLATA	CEL									X
PINNULARIA BREBISSONII VAR. RECTANGULARIS	CEL									X
PINNULARIA BREBISSONII VAR. STAURONEIFORMIS	CEL									X
PINNULARIA BREBISSONII VAR. TRINODIS	CEL									X
PINNULARIA BREBISSONII VAR. UNDULATA	CEL									X
PINNULARIA BREBISSONII VAR. VIRIDIS	CEL									X
PINNULARIA BRYOPHILA	CEL									X
PINNULARIA BRYOPHILA VAR. AMERICANA	CEL									X
PINNULARIA BRYOPHILA VAR. COMMUTATA	CEL									X
PINNULARIA BRYOPHILA VAR. GRUNOWII	CEL									X
PINNULARIA BRYOPHILA VAR. INTERRUPTA	CEL									X
PINNULARIA BRYOPHILA VAR. LINEARIS	CEL									X
PINNULARIA BRYOPHILA VAR. SUBCAPITATA	CEL									X
PINNULARIA BRYOPHILA VAR. UNDULATA	CEL									X
PINNULARIA BRYOPHILA VAR. VIRIDIS	CEL									X
PINNULARIA GIBBA	CEL									X
PINNULARIA GIBBA VAR. AMERICANA	CEL									X
PINNULARIA GIBBA VAR. COMMUTATA	CEL									X
PINNULARIA GIBBA VAR. GRUNOWII	CEL									X
PINNULARIA GIBBA VAR. INTERRUPTA	CEL									X
PINNULARIA GIBBA VAR. LINEARIS	CEL									X
PINNULARIA GIBBA VAR. SUBCAPITATA	CEL									X
PINNULARIA GIBBA VAR. UNDULATA	CEL									X
PINNULARIA GIBBA VAR. VIRIDIS	CEL									X
PINNULARIA MICROSTAURON	CEL									X
PINNULARIA MICROSTAURON VAR. AMERICANA	CEL									X
PINNULARIA MICROSTAURON VAR. COMMUTATA	CEL									X
PINNULARIA MICROSTAURON VAR. GRUNOWII	CEL									X
PINNULARIA MICROSTAURON VAR. INTERRUPTA	CEL									X
PINNULARIA MICROSTAURON VAR. LINEARIS	CEL									X
PINNULARIA MICROSTAURON VAR. SUBCAPITATA	CEL									X
PINNULARIA MICROSTAURON VAR. UNDULATA	CEL									X
PINNULARIA MICROSTAURON VAR. VIRIDIS	CEL									X
RHOICOSPHENIA CURVATA	CEL									X
SCHROEDERIA SETIGERA	CEL									X
SYNEDRA	CEL									X
SYNEDRA ULNA	CEL									X
TRACHELOMONAS	CEL									X
TRACHELOMONAS VOLVOCINA V. PUNCTATA	CEL									X
TOTAL				9005			1717			295

LAKE NAME: WILSON SINK RES.
STORET NUMBER: 3210

NYGAARD TROPHIC STATE INDICES

DATE	05 31 75	08 01 75	11 06 75
MYXOPHYCEAN	01/0 E	01/0 E	1.00 E
CHLOROPHYCEAN	0/0 O	03/0 E	0.33 ?
EUGLENOPHYTE	1.00 E	0/04 ?	0.25 E
DIATOM	0/06 ?	0/0 ?	0/07 ?
COMPOUND	02/0 E	04/0 E	1.67 E

PALMER'S ORGANIC POLLUTION INDICES

DATE	05 31 75	08 01 75	11 06 75
GENUS	05	00	05
SPECIES	03	00	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		05 31 75	08 01 75	11 06 75
AVERAGE DIVERSITY	H	1.47	1.66	2.57
NUMBER OF TAXA	S	10.00	7.00	19.00
NUMBER OF SAMPLES COMPOSITED	M	1.00	1.00	3.00
MAXIMUM DIVERSITY	MAXH	3.32	2.81	4.25
MINIMUM DIVERSITY	MINH	0.05	0.16	0.14
TOTAL DIVERSITY	D	3739.68	627.48	4055.46
TOTAL NUMBER OF INDIVIDUALS/ML	N	2544.00	378.00	1578.00
EVENNESS COMPONENT	J	0.44	0.59	0.60
RELATIVE EVENNESS	RJ	0.44	0.57	0.60
MEAN NUMBER OF INDIVIDUALS/TAXA	L	254.40	54.00	83.05
NUMBER/ML OF MOST ABUNDANT TAXON	K	1609.00	210.00	446.00

LAKE NAME: WILSON SINK RES,
STORET NUMBER: 3910

CONTINUED

TAXA	FORM	05 31 75			09 01 75			11 06 75		
		1B	%C	ALGAL UNITS PER ML	1B	%C	ALGAL UNITS PER ML	1B	%C	ALGAL UNITS PER ML
ANABAENA	FIL						X			X
APHANIZOMENON FLOS-AQUAE	FIL							11	8,7	137
CERATIUM HIRUNDINELLA	CEL						X			
CHROOMONAS ? ACUTA	CEL	13	24,5	623				14	21,7	343
CLOSTERIUM	CEL									X
COCCONEIS PLACENTULA	CEL							10	6,5	103
COELASTRUM MICROPORUM	COL									X
COSMARIUM	CEL								2,2	34
CRYPTOMONAS EROSA	CEL	11	63,2	1609						X
CRYPTOMONAS MARSSONII	CEL									X
CYMBELLA	CEL									X
CYMBELLA AFFINIS	CEL			X						
EPITHEMIA	CEL								4,4	69
EUGLENA	CEL			X						X
FLAGELLATES	CEL				12	55,6	210			
FRAGILARIA	CEL							12	33,9	377
GOMPHONEMA OLIVACEUM	CEL			X						
GYMNODINIUM	CEL				11	22,2	84			
HOUGEOTIA	FIL									X
NITZSCHIA #1	CEL	10	4,1	104						
NITZSCHIA #2	CEL	14	6,1	156						
OPEPHORA	CEL			X						
OSCILLATORIA	FIL			X					4,4	69
PENNATE DIATOM	CEL							13	28,3	446
RHOPALODIA GIBBA	CEL									X
SCENEDESMUS	COL				14	11,1	42			
SCHROEDERIA SETIGERA	CEL						X			
STAURASTRUM	CEL									
SYNEDRA ULNA	CEL	13	2,0	83						X
TETRALLRON MINIMUN V. SCROBICULATUN	CEL				13	11,1	42			
TOTAL				2544			378			1578

LAKE NAME: WALKER LAKE
STORET NUMBER: 3211

NYGAARD TROPHIC STATE INDICES

DATE	03 17 75	07 11 75	11 06 75
MYXOPHYCEAN	0/0 O	02/0 E	0/01 O
CHLOROPHYCEAN	01/0 E	0/0 O	0/01 O
EUGLENOPHYTE	0/01 ?	0/02 ?	0/0 ?
DIATOM	0/01 ?	0/0 ?	0/01 ?
COMPOUND	01/0 E	02/0 E	0/01 O

PALMER'S ORGANIC POLLUTION INDICES

DATE	03 17 75	07 11 75	11 06 75
GENUS	00	00	00
SPECIES	00	00	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE		03 17 75	07 11 75	11 06 75
AVERAGE DIVERSITY	H	1,34	0,01	0,03
NUMBER OF TAXA	S	3,00	2,00	2,00
NUMBER OF SAMPLES COMPOSITED	N	3,00	3,00	3,00
MAXIMUM DIVERSITY	MAXH	1,58	1,00	1,00
MINIMUM DIVERSITY	MINH	0,29	0,44	0,53
TOTAL DIVERSITY	D	64,32	0,10	0,09
TOTAL NUMBER OF INDIVIDUALS/ML	M	48,00	10,00	3,00
EVENNESS COMPONENT	J	0,85	0,01	0,03
RELATIVE EVENNESS	RJ	0,82	-0,76	-1,06
MEAN NUMBER OF INDIVIDUALS/TAXA	L	16,00	5,00	1,50
NUMBER/ML OF MOST ABUNDANT TAXON	K	27,00	10,00	3,00

LAKE NAME: WALKER LAKE
STORET NUMBER: 3211

CONTINUED

TAXA	FORM	03 17 75 IS	03 17 75 %C	03 17 75 ALGAL UNITS PER ML	07 11 75 IS	07 11 75 %C	07 11 75 ALGAL UNITS PER ML	11 06 75 IS	11 06 75 %C	11 06 75 ALGAL UNITS PER ML
CHROOCOCCUS	COL				1100.		10			
CHROOMONAS ?	CEL	133.31		16						
NAVICULA	CEL							100.		3
NODULARIA	FIL						X			
PINNATE DIATOM	CEL	166.31		27						
STAURASTRUM	CEL									X
TETRAEDRON MINIMUM	CEL	110.41		5						
TOTAL				48			10			3